

# BORON 15

## CONTAINS

Boron (B): 150 g/l

## RATES OF USE

Marginal Deficiency: 1.25 l/ha in at least 200 litres of water

Moderate Deficiency: 2.5 l/ha in at least 200 litres of water

Severe Deficiency: 2.5 l/ha in at least 200 litres of water initially followed by a further application after 3 – 4 weeks and again if symptoms should re-appear.

## TIMING

Liquibor can be applied to crops whenever a boron deficiency is observed or expected, after the 3 leaf stage, or when there is sufficient foliage present to absorb the spray.

Do not apply this product in very bright sunlight or in temperatures above 25°C. If possible, spray in the evening or early morning for best results.

## APPLICATION

**Shake the container well before opening and use.**

Spray equipment should be clean.

Half fill the spray tank with water, begin agitation and add the required quantity of Liquibor. Rinse the container thoroughly, add the rinsings to the spray-tank, complete the filling and apply without delay. Maintain agitation throughout.

Clean sprayer thoroughly after use.

Guard 2000® can be added at 0.1 %v/v of water volume to improve spray retention and rainfastness.

## COMPATIBILITY

Liquibor is physically and chemically compatible with a wide range of agrochemical products. Consult your supplier for the latest information. Up-to-date information is available at [www.headland-ag.co.uk](http://www.headland-ag.co.uk). Always read the label of the partner product carefully before mixing with Liquibor.

**STORAGE** Protect from frost. Store above 5°C. Keep in cool, well ventilated suitable store. Do not store in direct sunlight or near direct sources of heat. Keep out of reach of children. Keep away from food, drink and animal feedstuffs.

**SAFETY PRECAUTIONS** The product should not be considered as dangerous according to CLP (REG EC 1272/2008).

**Contents: 20 Litres**

# BORON 15